

EV0245

ePoster Viewing

Antimicrobials: resistance surveillance

Coagulase-negative staphylococci: pathogens in blood cultures and catheter-related infections, species distribution and susceptibility to linezolid, teicoplanin and vancomycin

S. Baka¹, I. Tsouma¹, S. Demeridou¹, V. Gennimata¹, E. Kouskouni¹

¹Aretaieio University Hospital, Athens, Greece

Objectives: Nosocomial bloodstream infections originating from central venous catheters (CVC) continue to represent a significant problem for hospitalized patients. By definition, derived from clinical studies, catheter-related bloodstream infections (CRBSI) are characterized by the isolation of the same microorganism (determined by antibiogram or DNA typing) from both blood and CVC cultures without evidence of infection with the identical pathogen at a different site. Although the most frequent causative microorganisms for CRBSI are coagulase-negative staphylococci (CNS), their clinical relevance is usually questionable. However, during the last decades, CNS have emerged as important nosocomial pathogens exhibiting decreased susceptibility to antimicrobials. The aim of this study was to assess the incidence of CRBSI caused by CNS and to determine the susceptibility to linezolid, teicoplanin and vancomycin of these isolates.

Methods: All hospitalized surgical patients at the Aretaieio University Hospital receiving a CVC between January 1st 2008 and December 31st 2013 were included in this study. Catheters were inserted by anesthetists in the operating theatre or post-anesthesia care unit or by intensive care physicians in the ICU. The catheter tip (about 5 cm) was cultured using the Maki method, by rolling it across a blood agar plate. After a 24 hours incubation period at 37°C, results were reported for growth equal to or exceeding 15 CFU/mL. The isolated pathogens were compared to the blood cultures' isolates. Blood cultures were performed using the automated BD BACTEC 9050 System. The identification as well as the susceptibility testing to the different antimicrobials of the isolated pathogens was performed using the automated system VITEK 2 (BioMérieux, Marcy l'Etoile, France).

Results: A total of 4,383 blood cultures and 356 catheter tips were screened during the period studied. Positive cultures with the same CNS from both catheter and blood culture were obtained in 119 cases. The CNS isolates were as follows: 78 (65.5%) *Staphylococcus epidermidis*, 22 (18.5%) *S. haemolyticus*, 14 (11.8%) *S. hominis*, 3 (2.5%) *S. saprophyticus* and 2 (1.7%) *S. capitis*. All isolates were susceptible to linezolid, while 5 (6.4%) *S. epidermidis*, 1 (7.1%) *S. hominis* and 1 (4.5%) *S. haemolyticus* isolates were resistant to teicoplanin. Finally, 2 (2.6%) *S. epidermidis* isolates were resistant to vancomycin.

Conclusion: CNS are a group of microorganisms that are increasingly implicated in a plethora of infections, including device-related. The most frequent species of CNS isolates were *S. epidermidis*. All isolates were susceptible to linezolid and only few were resistant to glycopeptides. In order to reduce the incidence of CRBSI it is imperative to adhere to evidence-based practice when placing and maintaining CVC. CNS resistant strains remain a major problem since they can serve as a reservoir of resistance genes.