

P1681

Abstract (poster session)

Carbapenemases arrived in Germany: report for 2011 of the National Reference Laboratory for Multidrug-Resistant Gram-negative Bacteria

M. Kaase*, F. Szabados, A. Anders, S. Gatermann (Bochum, DE)

Objectives: Multidrug-resistance in Enterobacteriaceae, *Pseudomonas aeruginosa* and *Acinetobacter baumannii* is of utmost therapeutic importance since no innovative antimicrobial drugs against gram-negative bacteria will be introduced into clinical practice within the next five years. Among all resistance mechanisms the worldwide spread of carbapenemases is the most worrisome development. However, the correct identification of carbapenemases is challenging for the microbiological laboratory. **Methods:** The National Reference Laboratory for Multidrug-Resistant Gram-negative Bacteria offers the free service of carbapenemase detection in bacterial isolates with elevated carbapenem MICs. All isolates are tested by a wide array of phenotypic and molecular methods. A bioassay based on cell-free extracts allows the detection of still unknown beta-lactamases. **Results:** A total of 1074 isolates were sent to the National Reference Laboratory in 2011 between January 1st and October 11th mainly for investigation for carbapenemases, but also for clarification of the resistance mechanism to 3rd generation cephalosporins or molecular strain typing. Several different carbapenemases could be detected, including OXA-48 (n = 79), OXA-162 (n = 4), OXA-181 (n = 1), OXA-204 (n = 1), KPC-2 (n = 41), KPC-3 (n = 9), VIM-1 (n = 47), VIM-2 (n = 28), VIM-4 (n = 4), VIM-26 (n = 1), IMP-7 (n = 1), IMP-8 (n = 7), IMP-13 (n = 2), IMP-31 (n = 1), NDM-1 (n = 16), GIM-1 (n = 2), OXA-23 (n = 134), OXA-72 (n = 7) and OXA-58 (n = 2). In Enterobacteriaceae most Carbapenemases were found in *K. pneumoniae*, especially OXA-48, KPC-2 and KPC-3. VIM-2 was the most frequent carbapenemase in *P. aeruginosa* and OXA-23 in *A. baumannii*. **Conclusion:** Almost all carbapenemases found worldwide have arrived in Germany. However, the molecular epidemiology in Germany with a predominance of OXA-48 differs significantly from observations made in other countries like Greece, Israel, USA or the United Kingdom. An ongoing surveillance of resistance determinants is necessary, especially for infection control and diagnostics.