

Session: P070 Update on respiratory viruses

**Category: 1c. Influenza and respiratory viruses**

24 April 2017, 13:30 - 14:30  
P1401

**The first report of detection and phylogenetic analysis of enterovirus D-68 and rhinoviruses in children under age of 5 years in Iran**

Nastaran Ghavami<sup>\*1</sup>, Haideh Namdari<sup>2</sup>, Farhad Rezaei<sup>3</sup>

<sup>1</sup>*Tehran University of Medical Sciences; School of Public Health, Virology Department*

<sup>2</sup>*Tehran University of Medical Sciences; Immunology*

<sup>3</sup>*Tehran University of Medical Sciences; Virology Department*

**Background:** Enterovirus D68 (EV-D68) was reported to cause an outbreak of severe respiratory illness in children needing hospital or critical care admission across the United States in 2014. Also, Rhinoviruses are the most common cause of respiratory tract infections in individuals of all ages, causing one-half of common colds annually and 90% of colds during the autumn epidemic season. The goal of this study was to determine frequency and genotype characterization of EV-68 and Rhinoviruses in children under age of 5 years for the first time in Iran.

**Material/methods:** A total of 291 children hospitalized for any respiratory infection between September, 2015 and April, 2016 were screened for EV-D68 and rhinoviruses. The presence and quantity of EV-D68 and rhinoviruses in nasal swab samples were determined by Real-Time RT-PCR and nested RT-PCR methods. Positive samples were assessed further for genotype characterization by partial sequence analysis and phylogenetic analysis.

**Results:** A total of 15/291 (5.16%) of samples tested positive for EV-D68, and 61/291 (20.96%) cases were positive for rhinoviruses. Mean age was 4.3 and 3.8 years in the EV-D68 and rhinoviruses positive groups, respectively. Children with EV-D68 infection were more likely to have difficulty breathing. There was no significant difference in admission to the critical care unit among children with EV-D68 infection compared with those with rhinovirus infections. Children with EV-D68 infection were more often admitted to hospital, but not significantly. Children positive for EV-D68 or rhinoviruses type C were had same rate of wheezing than rhinoviruses type A and B (*odds ratio*; 2.73 and *pvalue*; 0.04).

**Conclusions:** Rhinoviruses are more frequently detected from children than EV-D68. However, Enterovirus D68 seems to be a more virulent pulmonary pathogen than rhinoviruses, but we could not find a significant difference in need for critical care.