

Erasmus MC
Centrum

Pk/Pd of antifungal drugs

Johan W. Mouton MD PhD FIDSA
Professor pharmacokinetics and pharmacodynamics

JWM New Delhi 19-03-2015 Radboud Universiteit Nijmegen Radboudumc

Disclosures

Erasmus MC
Centrum




Research grants – advisory boards – speaker

JWM New Delhi 19-03-2015 Radboud Universiteit Nijmegen Radboudumc

Erasmus MC
Centrum


Dosing should be such that the level of antimicrobial activity is associated with a high likelihood of therapeutic success.





therapeutic success


DOSE


JWM New Delhi 19-03-2015 Radboud Universiteit Nijmegen Radboudumc





























EUCAST EUROPEAN COMMITTEE ON ANTIMICROBIAL SUSCEPTIBILITY TESTING

Susceptible (S)

A micro-organism is defined as susceptible by a level of antimicrobial activity associated with a high likelihood of therapeutic success. A micro-organism is categorized as susceptible by applying the appropriate breakpoint in a defined phenotypic test system.

Note: This breakpoint may be altered with legitimate changes in circumstances

Intermediate (I)


A micro-organism is defined as intermediate by a level of antimicrobial activity associated with indeterminate therapeutic effect. A micro-organism is categorized as intermediate by applying the appropriate breakpoints in a defined phenotypic test system.

Note: This breakpoint may be altered with legitimate changes in circumstances.

Resistant (R)

Bacteria are defined as resistant by a level of antimicrobial activity associated with a high likelihood of therapeutic failure. A micro-organism is categorized as resistant by applying the appropriate breakpoint in a defined phenotypic test system.

Note: This breakpoint may be altered with legitimate changes in circumstances





Problem (or is it?):


What if the standard dose is different?


What if the population is different?

JWM New Delhi 19-03-2015









Pharmacokinetic parameters :
Measures of Exposure

AUC is *usually* linearly related to Dose

Dose x 2 = AUC x 2
Dose x 4 = AUC x 4

Erasmus MC
JWM New Delhi 19-03-2015 Radboud Universiteit Nijmegen Radboudumc


So what determines the relationship between dose and exposure?

Erasmus MC
JWM New Delhi 19-03-2015 Radboud Universiteit Nijmegen Radboudumc

Fluconazole Dose - AUC Relationship

$r=0.99$

Rodriguez-Tudela et al, AAC 2007
Erasmus MC
JWM New Delhi 19-03-2015 Radboud Universiteit Nijmegen Radboudumc





Erasmus MC
Centrum


So what more determines the relationship between dose and exposure?


JWM New Delhi 19-03-2015 Radboud Universiteit Nijmegen Radboudumc

















Erasmus MC
Centrum

EUCAST

EUROPEAN COMMITTEE
ON ANTIMICROBIAL
SUSCEPTIBILITY TESTING

Susceptible (S)

A micro-organism is defined as susceptible by a level of antimicrobial activity associated with a high likelihood of therapeutic success. A micro-organism is categorized as susceptible by applying the appropriate breakpoint in a defined phenotypic test system.

Note: This breakpoint may be altered with legitimate changes in circumstances

Intermediate (I)

A micro-organism is defined as intermediate by a level of antimicrobial activity associated with intermediate therapeutic effect. A micro-organism is categorized as intermediate by applying the appropriate breakpoints in a defined phenotypic test system.
Note: This breakpoint may be altered with legitimate changes in circumstances.

Resistant (R)

Bacteria are defined as resistant by a level of antimicrobial activity associated with a high likelihood of therapeutic failure. A micro-organism is categorized as resistant by applying the appropriate breakpoint in a defined phenotypic test system.
Note: This breakpoint may be altered with legitimate changes in circumstances.

JWM New Data 19-03-2015 Radboud Universiteit Nijmegen Radboudumc

LAB REPORT

Erasmus MC
Centrum

Sensitivity	Organism 1	Escherichia coli
Hoeveelheid		>=10E5 kve/ml
Panel gevoeligheid		5 Urine Coliform
amoxicilline/clavula		Sensitive (0,06 mg/l)
amoxicilline		Sensitive (0,06 mg/l)
cefuroxim		Sensitive (0,06 mg/l)
cefotaxim		Sensitive (0,5 mg/l)
cefazoline		Sensitive (0,25 mg/l)
ciprofloxacine		Sensitive (<+0,06 mg/l)
doxycycline		Sensitive (1 mg/l)
nitrofurantoin		Sensitive (<+32 mg/l)
norfloxacine		Intermediate (1 mg/l)
sulfamethoxazol		Sensitive (<+64 mg/l)
tobramycine		Intermediate (0,25 mg/l)
trimethopim		Resistant (>64 mg/l)
cotrimoxazole		Sensitive (1 mg/l)
ceftazidim		Sensitive (0,13 mg/l)

- Provides Clinician/Consultant guidelines how to optimally treat a patient (Freely translated from EUCAST guideline)

BASED ON EXPOSURES OF COMMON DOSES

JWM New Data 19-03-2015 Radboud Universiteit Nijmegen Radboudumc

LAB REPORT

Erasmus MC
Centrum

Sensitivity	Organism 1	Escherichia coli
Hoeveelheid		>=10E5 kve/ml
Panel gevoeligheid		5 Urine Coliform
amoxicilline/clavula		Sensitive (0,06 mg/l)
amoxicilline		Sensitive (0,06 mg/l)
cefuroxim		Sensitive (0,06 mg/l)
cefotaxim		Sensitive (0,5 mg/l)
cefazoline		Sensitive (0,25 mg/l)
ciprofloxacine		Sensitive (<+0,06 mg/l)
doxycycline		Sensitive (1 mg/l)
nitrofurantoin		Sensitive (<+32 mg/l)
norfloxacine		Intermediate (1 mg/l)
sulfamethoxazol		Sensitive (<+64 mg/l)
tobramycine		Intermediate (0,25 mg/l)
trimethopim		Resistant (>64 mg/l)
cotrimoxazole		Sensitive (1 mg/l)
ceftazidim		Sensitive (0,13 mg/l)


- Provides Clinician/Consultant guidelines how to optimally treat a patient (Freely translated from EUCAST guideline)

BASED ON EXPOSURES OF COMMON DOSES IN ADULTS

JWM New Data 19-03-2015 Radboud Universiteit Nijmegen Radboudumc

Erasmus MC
Centrum

Exposures in children : differences




- Lower EXPOSURE then expected
- Clearance in individual

JWM New Delhi 19-03-2015 Radboud Universiteit Nijmegen Radboudumc

Erasmus MC
Centrum

Pharmacokinetic parameters of Fluconazole by age group




Mouton, Antimicrob Pharmacodynamics in Theory and Clin Practice 2007, 357

JWM New Delhi 19-03-2015 Radboud Universiteit Nijmegen Radboudumc


Erasmus MC
Centrum


AUC per unit dose (mg/kg) of fluconazole differences by age group




Mouton, Antimicrob Pharmacodynamics in Theory and Clin Practice 2007, 357

JWM New Delhi 19-03-2015 Radboud Universiteit Nijmegen Radboudumc







Erasmus MC
Centrum

Exposures in children:

- Generally lower – increase dose
- Larger variability – be aware, possible increase dose *or* monitor *or* estimate clearance
- These insights can help to rationalize dosing regimens in children

JWM New Data 19-03-2015 Radboud Universiteit Nijmegen Radboudumc

Erasmus MC
Centrum

To summarize:

- There is a good relationship for exposure and response
- This translates to dosing regimens *for the general population*
- Breakpoints are based on *the most common lowest dose*
- Other populations may require dose adjustments
- For non-predictable concentrations: TDM is a requirement

JWM New Data 19-03-2015 Radboud Universiteit Nijmegen Radboudumc

ESCMID Online Lecture Library © by author