

P1519

Paper Poster Session

Antibacterial resistance: the animal and food production part of one health

Prevalence of ESBL and/or carbapenemase-producing *Escherichia coli* isolated from yellow-legged gulls from Barcelona, Spain

Andrea Vergara Gómez*¹, Cristina Pitart², Tomás Montalvo³, Juan Carlos Hurtado¹, Sara Sabate⁴, Raquel Planell⁴, Francesc Marco Reverte¹, Beatriz Ramírez⁴, Victor Peracho⁴, Mercé de Simón⁴, Jordi Vila¹

¹*Department of Clinical Microbiology-Cdb, Hospital Clínic - Universitat de Barcelona, Barcelona, Spain*

²*Department of Clinical Microbiology-Cdb, Hospital Clínic - Universitat de Barcelona; Institut de Salut Global (Isglobal-Hospital Clínic), , Barcelona, Spain*

³*Agencia de Salut Pública de Barcelona, Barcelona , Spain*

⁴*Agencia de Salut Pública de Barcelona, Barcelona, Spain*

Background: In the last decade, the number of bacterial pathogens presenting multidrug resistance to antibacterial agents has increased dramatically, becoming an emergent global disease and a major public health problem. The main cause behind the increasing rates of resistance can ultimately be found in the abuse and misuse of antibacterial agents, whether used in patients and livestock or released into the environment. Once antimicrobial resistant bacteria emerge, they can spread locally or globally. The main factors contributing to their spread at a global level comprise migrant birds, globalization of commercial food and international traveling. The objective of this study was to investigate the prevalence of ESBL and/or carbapenemase-producing *Enterobacteriaceae* from faecal swabs obtained from a group of yellow-legged gulls in Barcelona, Spain.

Material/methods: One hundred and thirty-two faecal swabs from yellow-legged gulls were obtained during a one-month period. The samples were plated on ESBL chromogenic agar. Identification of ESBL-producing enterobacteria was carried out by mass spectrometry (MALDI-TOF). Antimicrobial susceptibility was tested by disk diffusion. Characterization of ESBL and carbapenemases genes was performed by PCR followed by sequencing (ESBL: *bla*_{TEM}, *bla*_{SHV}, *bla*_{CTX-M}; carbapenemases: *bla*_{KPC}, *bla*_{OXA-48}, *bla*_{VIM}, *bla*_{IMP} and *bla*_{NDM}; cephamycinases: *bla*_{CMY}, *bla*_{DHA}, *bla*_{FOX}, *bla*_{ACC}, *bla*_{EBC}, *bla*_{MOX}). To evaluate the genetic relationship among the strains, REP-PCR was performed. Agglutination with antiserum O25 was used to know if those CTXM-15-carrying isolates belong to the high risk clone O:25b-ST131.

Results: Seventy two (54%) out of 132 faecal samples were positive for ESBL, carbapenemases or cephamycinases producing *Escherichia coli* (Table 1). Forty-five strains (62.5%) were resistant to quinolones, 22 (30.6%) to gentamicin and 9 (12.5%) to amikacin. Rep-PCR showed a high genetic heterogeneity among the strains. Those strains with CTXM-15 did not show agglutination with

antiserum O25. The two isolates carrying the KPC and VIM carbapenemases were not genetically related and belonged to the ST1011 (phylogroup E) and ST354 (phylogroup F), which have been previously found in humans.

Conclusions: The yellow-legged gulls screened in the present study presented a high prevalence of ESBL-producing *E. coli*. The isolation of two carbapenemase-producing *E. coli* strains is a matter of concern. These results reinforce the knowledge that the feeding and migratory habits of these birds play an important role in the dissemination of antibiotic-resistant bacteria between geographically distant ecosystems.

Table 1. Distribution of β -lactamases.

β -lactamase	N° strains	%
SHV Group	38	52.8
SHV-12	24	33.3
SHV-12 + TEM-1	13	18
SHV-2	1	1.4
CTX-M group	30	41.6
CTX-M-15	11	15.3
CTX-M-15 + TEM-1	2	2.8
CTX-M-1	1	1.4
CTX-M-1 + TEM-1	5	6.9
CTX-M-1 + TEM-84	1	1.4
CTX-M-14	4	5.5

CTX-M-14 + TEM-1	6	8.3
VIM-1 + KPC-2	2	2.8
CMY-2	2	2.8
TOTAL	72	100