

# BONE AND JOINT INFECTIONS (BJI) SINCE LABELLING OF COMPLEX BJI REFERENCE CENTRES IN FRANCE: BEFORE/AFTER STUDY

Laurent E.<sup>1,2,3,4</sup>, Gras G.<sup>2,5</sup>, Druon J.<sup>2,6</sup>, Rosset P.<sup>2,6</sup>, Baron S.<sup>1,2,3</sup>, Le Louarn A.<sup>1,2</sup>, Rusch E.<sup>1,2,3,4</sup>, Bernard L.<sup>2,4,5</sup>, Grammatico-Guillon L.<sup>1,2,3,4</sup>

<sup>1</sup> Epidemiology Unit; <sup>2</sup> University Hospital of Tours, France; <sup>3</sup> Research Team EE1 EES; <sup>4</sup> F. Rabelais University, Tours, France; <sup>5</sup> Infectious Diseases Unit; <sup>6</sup> Orthopaedic Unit

## Introduction

- BJI are associated with morbimortality and high economic impact
- First BJI national study through French Hospital Discharge Database (HDD) in 2008:
  - validation of a BJI case definition
  - 1<sup>st</sup> prevalence estimation: 54/100,000 inhabitants<sup>1</sup>

## Objectives

- Epidemiologic and medico-economic evolution of BJI between 2008 and 2013
- Focus on CBJI in 2013

## Methods

- Extraction of BJI stays of patients ≥15 from French national HDD
- Identification of native and device-associated BJI (DBJI) with the validated algorithm using specific:
  - ICM-10 diagnosis codes
  - surgical procedure codes according to the French Common Classification of Medical Acts<sup>1</sup>
- CBJI Identification: Z76800 code:
  - created in 2011
  - coded after a multidisciplinary consultation in FRC
- Stay HD → patient HD (linkage with the unique and encrypted patient number)

➤ BJI Prevalence between 2008 and 2013, from 54/100,000 to 70/100,000 inhabitants (37,252 inpatients)

Table

Hospital stays	Overall BJI			Device-associated BJI			Complex BJI	
	2008		2013		2008		2013	
	N	%	N	%	N	%	N	%
Hospital stays	36,097	43	48,386	57	11,451	32**	16,214	34**
Public sector hospitalization	28,441	78.8	39,797	82.2	7,923	69.2	12,102	74.6
FRC hospitalization	11,170	30.9	15,888	32.8	3,448	30.1	5,448	33.6
Surgical stay wards	19,847	55.0	27,308	56.4	8,513	74.3	12,087	74.5
Resuscitation/ICU stays*	2,106	5.8	3,350	6.9	624	5.4	991	6.1
Z76800 code (2013 only)	-	-	3,301	6.8	-	-	2,033	12.5
Device-associated BJI	11,451	31.7	16,214	33.5			2,033	61.6
Microorganisms	14,407	39.9	30,833	63.7	4,708	41.1	11,077	68.3
Bacteria	14,071	97.7	30,311	98.3	4,668	99.2	10,998	99.3
Polybacteria	1,591	11.0	6,116	19.8	502	10.7	1,716	15.5
Staphylococci	9,425	65.4	20,837	67.6	3,570	75.8	8,484	76.6
Streptococci	2,208	15.3	5,678	18.4	687	14.6	1,636	14.8
Gram negative Bacilli	2,370	16.5	8,104	26.3	788	16.7	2,373	21.4
Resistance	485	3.4	7,974	25.9	183	3.9	2,925	26.4
LOS (days), mean (median [min-max])		17.5	17.5		18.9	18.2		22.2
	(11 [1-421])		(12 [1-442])		(13 [1-421])		(13 [1-386])	

\* Intensive Care Unit

\*\* % overall BJI

BJI (Table)

➤ Similarities 2008-2013:

- Type: 50% septic arthritis, 37% osteomyelitis, 10% spondylodiscitis
- ➤ prevalence with age and sex (SR M/F 1.5) (Fig. I)
- 1 comorbidity or more for 2/3 of inpatients (cardiac disease, diabetes, Fig. II)
- Hospital case-fatality rate 5%
- Rehospitalisations 20%
- Mean length of stay (LOS) 17.5 days

➤ Differences 2008-2013:

- ➤ coded microorganisms: 40 to 64%
- ➤ costs (Fig. III)

DBJI (Table)

➤ 34% of BJI, stable

➤ Microorganisms: Staphylococci 77%

➤ Differences 2008-2013:

- ➤ hospitalisations in public sector
- ➤ costs (Fig. III)
- ➤ mean LOS from 18.9 to 18.2 days ( $p<0.01$ )

## Results

Figure I

All BJI / DBJI Prevalence, France, 2008-2013





Figure II

BJI Comorbidities, France, 2008-2013



\* PVD : Peripheral Vascular Disease


## Discussion

- Increasing prevalence of BJI and DBJI.
- DBJI more frequent in men, elderly people, with comorbidities, as in 2008.
- Increasing prevalence could be due to:
  - Better knowledge + financial incentive since the labelling of FRC.
  - An ageing population with more comorbidities, undergoing more arthroplasties.
  - A global activity increase.
- Estimation of CBJI prevalence, required for FRC monitoring, complicated by:
  - the absence of clinical consensus about CBJI definition.
  - An underuse of the Z76800 code, recently created and needing a multidisciplinary consultation.
- Could CBJI epidemiology be approached by the study of DBJI ?

<sup>1</sup> Grammatico-Guillon L *et al.* Bone and joint infections in hospitalized patients in France, 2008: clinical and economic outcomes. *J. Hosp. Infect.* 82, 40-48 (2012)

Figure III

BJI costs (euros), France, 2008-2013



CBJI (Table)

➤ Z76800 code sparsely used: 7% of hospital stays

○ More used in DBJI (12.5%) than in native BJI (3.5%).

○ Used in 20% of BJI stays and 34 % of DBJI stays in FRC.