

E0143 *Acinetobacter baumannii* from a sewage of a nursing home in Croatia

Branka Bedenic*¹, Natasa Belder¹, Marko Siroglavic², Mia Slade³, Haris Car⁴, Svjetlana Dekic⁵, Martina Seruga Music^{5,5}, Aleksandra Presecki-Stanko⁶, Dorotea Šijak⁷, Jasna Hrenovic⁵

¹School of Medicine, University of Zagreb, Clinical Department for Clinical and Molecular Microbiology, Zagreb, Croatia, ²University Hospital Center Zagreb, Transfusiology, Zagreb, Croatia, ³Institute for Transfusiology, Transfusiology, Zagreb, Croatia, ⁴Secondary Medical School, Zagreb, Croatia, ⁵University of Zagreb, Faculty of Science, Biology, Zagreb, Croatia, ⁶University Hospital Center Zagreb, Clinical Department for Clinical and Molecular Microbiology, Zagreb, Croatia, ⁷School of Medicine, University of Zagreb, Zagreb, Croatia

Background: In the previous studies the predominance of OXA-24/40 and OXA-23 β -lactamase was reported among *Acinetobacter baumannii* in both the hospitals and long-term care facilities (LTCF) in Croatia. The aim of this study was to determine antibiotic susceptibility, mechanisms of antibiotic resistance and molecular epidemiology of *A. baumannii* isolates recovered from sewage of a nursing home in Zagreb.

Materials/methods: In April 2017, 10 *A. baumannii* were isolated from sewage. Antibiotics susceptibility was determined by broth microdilution method. The presence of carbapenemases of class A, B and D and extended-spectrum β -lactamases genes was explored by PCR. The occurrence of the IS*Aba1* upstream of the *bla*_{OXA-51-like} or *bla*_{OXA-23-like} was determined by PCR mapping. Conjugation and transformation experiments were performed as previously described. Plasmid incompatibility groups were determined by PBRT. Genotyping were performed by sequence group determination, PFGE and MLST.

Results: Eight isolates were resistant to gentamicin and ciprofloxacin, three to ceftazidime and cefepime and two to carbapenems. There was uniform susceptibility to colistin. All isolates were positive for chromosomal *bla*_{OXA-51} whereas two carbapenem-resistant isolates were found to possess also *bla*_{OXA-23} genes. IS*Aba1* was found upstream of *bla*_{OXA-51} in seven out of ten isolates. Attempts to transfer imipenem resistance were unsuccessful indicating chromosomal location of *bla*_{OXA-23} gene. The plasmids extracted from isolates positive for OXA-23 belonged to Inc group 6 encoding *aci6*-replicase gene.

Two OXA-23 positive isolates belonged to SG 1 (IC2) whereas the rest of the isolates susceptible to carbapenems were allocated to SG 2 (IC1). Additional MLST analysis corroborated that two OXA-23 and carbapenem-resistant isolates belong to the ST-195 clustering into the CC92 within the IC2, commonly reported worldwide. The susceptible isolates belonging to IC1 were found to be clonally related by PFGE, but different from the carbapenem-resistant .

Conclusions: This study showed dissemination of OXA-23 producing *A. baumannii* from the nursing home into the sewage. Disinfection of nursing home sewage should be recommended in order to prevent the spread of resistance genes into the community sewage and nature. In 2013, OXA-23 positive *A. baumannii* from investigated nursing home were found to belong to ST-487, demonstrating the emergence of a new clone.